



Return-to-Sport after ACL Revision Reconstruction Protocol

- Allograft or Hamstring Tendon Graft- Immobilizer and crutches for 2 weeks
 - Initial rehabilitation protocol should follow the same criteria and goals, however, progression to activities such as running, jumping (plyometrics), and cutting may be delayed depending on how the patient presents with recovery and healing.
 - o Delayed as much as 4-6 months, depending on the fixation on the soft tissue in the bone tunnels and individual patient healing time.
 - Healing time for the autograft hamstring (tendon-to-bone) is approximately 8-12 weeks
 - o *** No resistive Hamstring strengthening until weeks 6-8 to allow for graft site healing**
 - Revascularization of graft occurs at 8-12 weeks post-op- Rodeo et al. JBJS 1993
 - * Not a good candidate for an accelerated Rehabilitation Protocol or early running, jumping, cutting, etc. (2° to slower soft tissue->bone healing)**
 - Be aware of patellofemoral type pain, if it does arise, treat it like Patellofemoral Pain Syndrome with modalities, taping techniques, myofascial release techniques, NMES for quad strength regeneration
 - o Also evaluated the entire kinetic chain, look for dysfunctional patterns along the chain, especially in the hip, ankles, SI, and Lumbar spine
 - Kinetic Chain Ripple Effect- everything is connected
 - o Functional movement is through all 3 planes, and there is a mobility and stability component at each joint and with each muscle for proper movement
 - **Example: Gluteus Maximus**, classified as an extender of the hip, look at the alignment of the muscle fibers, just as much a hip abductor and external rotator. It acts as a prime mover and a stabilizer at the same time. Moving in one, while stabilizing in the other two, depending on the position of the joint and the movement the athlete is performing
 - All muscles/movements have to be train for both strength and stability
 - You need to look at soft tissue: connected fascially and interact for true functional movement
 - **Example: SLR= Strain on multiple tissues, not just the hamstring**
 - o ITB, ipsilateral lumbar fascia, Achilles tendon, contralateral lumbar fascia, and plantar fascia in descending percentages
- * This protocol is a criterion-based protocol, with a goal of maximizing healing and strength recovery, while minimizing risk of re-injury, so the athlete can return, as quickly and safely as possible, to athletic activities at the pace that each individual progresses.

Immediately Post-Operative Phase: (Week 1, approx. days 1-7 post-operatively)

Goals of this phase are:

- Restore full passive knee extension, and gradually ↑ knee flexion to 90°
- Diminish joint swelling and pain
- Restore patellar mobility
- Re-establish quadriceps control- active quadriceps contraction with superior patellar glide
- Improve ambulation to FWB/ ↓ assistive device use

Treatments: Perform/increase number and duration of exercises as tolerated

- Gait Training:
 - o Patient is WBAT in full locked extension brace with bilateral crutches for 2 weeks Post-op
- Exercises:
 - o Ankle Pumps to ↓ swelling/edema
 - o Ankle resistance band open chain exercises- all 4 directions

- Patellar mobilization in all directions
- Quadriceps isometric setting, with VMO self biofeedback
- Gluts sets
- Hamstring stretches for hamstring re-lengthening
- Straight Leg Raises (flexion, abduction, adduction)
- Gastrocnemius/heel cord stretching for re-lengthening
- Active and Passive knee flexion exercises to tolerance, not pushed- (to 90° by days 5-7)
- Wall slides (supine with foot on wall/window) for increasing knee flexion, or seated AAROM
- Gentle overpressure into full extension (PT or pt. actively)
- Weight Bearing: weight shifts in full extension brace
- NMES:
 - Can and should be used, if contraction deficit is present, during active muscle exercises
- CP with full Extension, could also use Kinesio Tape (basket weave technique) or dry needling technique, to control swelling/edema
- Patient Education: What to expect, Goals of the phases, contraindicated movements/activities, etc.-
Continued throughout all phases of the protocol
- Home Exercises Program- See HEP for Immediately Post-op Phase

Early Post-Operative Phase:

Criteria for entering this phase is:

- 1- Quadriceps voluntary initiation with quad set and SLR
- 2- Full passive knee extension
- 3- Knee A + PROM of 0°-90°
- 4- Good patellar mobility
- 5- Minimal joint effusion
- 6- Independent ambulation

Goals of this phase are:

- To maintain full passive and active knee extension
- To gradually increase knee flexion to 110°-120°
- To diminish swelling and pain
- Increase muscular training
- Restore proprioception
- Maintain patellar mobility
- Use a recumbent cycle without difficulty
- Perform a SLR with no extension lag
- Reciprocal stair climbing
- KOS-ADL Score of >65%

Treatment:

- Gait Training: **Immobilizer worn for 1-2 additional weeks.** Then WBAT, with a goal of discontinuing use of crutches by day 14-18 post-op- continue brace until voluntary quad control is demonstrated, ***May be subject to change by MD**
- Exercises: (**continue/progress all exercises from previous phase**)
 - Continue NMES with quadriceps exercises, if needed, Quad Sets
 - SLR in all 4 planes- progression with 1# weight (***Proximal loading**)
 - PROM 0° to 110°-120°, OKC –Knee Extension 90° to 40°
 - Mini Squats 0°-30° DL (BW only) Progress to unstable surface such as tilt board (Med./Lat. & Ant./Post.) or foam with 3-5 second holds ***Forward trunk tilt- recruit H/S & unload ACL**, Wall Squats/Sits to 90°
 - Overpressure into Full Knee Extension, Prone Knee Hangs if lacking full extension
 - Hamstring Curls

- Weight Shifts- Medial/Lateral & Diagonal Patterns, Balance Weight Shifting- balance board, BOSU
- Step-Ups in pain free range- Front & Lateral
- Lateral Step Over: Cones, Hurdles- ***Instruct to raise knee to level of hip**
- DL Leg Press- Sub Maximal
- Stationary Bicycle (if ROM permits)
- Manual Treatments:
 - Patellar Mobilization (If flexion is limited), Scar Mobs, (If skin is healed)
- Continue to control swelling: Ice, elevation if needed

Intermediate Post-Operative Phase:

Criteria for entering this phase:

- 1- P & AROM knee flexion to $\geq 110^\circ$
- 2- Quadriceps & Hamstring strength $\geq 60\%$ of the non-injured leg (Dynamometry)
- 3- Minimal to no effusion
- 4- No joint line or patellofemoral pain
- 5- Mild laxity on clinical examine, or +1 or less with the KT-2000 test (if available)
 - a. Week 4 & 6: KT-2000 test at 20 & 30 lbs anterior and posterior

Goals of this phase are:

- Quadriceps strength to $\geq 75\%$ of the non-injured leg, H/S $\geq 75\%$ non-injured leg
- Restore knee ROM to 0° - 125° , or knee ROM to within 10° - 15° of non-injured leg
- Improve lower extremity strength
- Enhance proprioception, balance, and neuromuscular control
- Improve muscular endurance
- Restore limb confidence and function
- KOS-ADL score or $>75\%$
- Get an Post-Op score for IKDC or KOOS (age appropriate version)

Treatments:

- Gait Training: Unlocked brace, discontinued use if sufficient quad strength is present. Retrain with normal walking gait pattern
- Exercises: (**Continue/progress all exercises from previous phases as tolerance/as needed**)
 - Knee concentric & eccentric $90^\circ \leftrightarrow 40^\circ$ -(no weights) at 4-5 weeks, isometrics can be begun at 6-7 weeks post-op
 - Front & Lateral step down exercises, Progress Lateral step-up exercise
 - Heel raises (raising up onto toes)
 - Forward & Lateral Mini Lunges (30° - 40° flexion)- **if good quad control**
 - Wall Squats (0° - 30°)- **if patient has good quadriceps control**
 - Perturbation training – DL-> progress to →SL Balance when appropriate- Can also use uneven surfaces and external perturbations to progress (DL & SL)- Tilt Board (stabilized to level position initially), BOSU®, Airex®- progress with ball toss & catch/reaching (UE & LE), external tapping on hips and trunk on any surface- ***With knee slightly flexed**
 - Pool exercises if available: forward & backward walk/run (***forward at week ≥ 6**), hip & leg exercises- at slow speeds of mvmt. (Graft protection)
 - Begin testing/training for proprioception on the Biodex® Stability System (or other system)- ***if available, or:** Balance Error Scoring System Test, Functional Reach Test, Tandem Walking tests- See attached sheets
 - Progress bike to minimum of 10 minutes for increasing endurance
 - Core stabilization/strengthening- Bridging (Progress with ball squeeze→ unstable surface)
- Manual Treatments:
 - Continue patellar and scar mobilization if needed
 - Tibiofemoral mobilization with rotation for ROM if joint mobility is limited.

Late Post-Operative Phase:

Criteria for entering this phase:

- 1- Active knee ROM 0° - $\geq 125^{\circ}$
- 2- Quadriceps Strength & Girth $\geq 75\%$ of non-injured leg (dynamometry & tape measure), Hamstring Strength $\geq 75\%$ of non-injured leg (dynamometry)

Goals for this phase:

- Restore full knee ROM
- Continue to improve lower extremity strength, balance, proprioception, neuromuscular control, muscular endurance
- Knee effusion to trace or less
- Quadriceps strength to $>80\%$, Hamstring Strength $>80\%$ of the non-injured leg Normal gait pattern
- Minimal laxity on clinical exam, or 2mm or less on the KT-2000 test
- Get a baseline SFMA Assessment at ≥ 10 weeks post-op

Treatment:

- Exercises: (**Continue/progress all exercises from previous phases as tolerance/as needed**)
 - o Progress exercises in intensity and duration
 - o Light isotonic hamstring exercises begun 7-8 weeks post-op
 - o Perturbation training continues
 - o Mini Squats 0° - 30° DL (BW only) Progress to unstable surface such as tilt board (Med./Lat. & Ant./Post.) or foam with 3-5 second holds ***Forward trunk tilt- recruit H/S & unload ACL**, Progress to 45° - 60°
 - o Dynamic/Plyometric Leg Press (Begin at ≥ 10 Weeks- To learn technique and control ground reaction forces, "land softly on toes with knees slightly flexed" for dissipation of force and to avoid hyperextension)
 - o Progress lateral stepping and lateral step-down exercises with resistance bands on the distal femur creating a medial pull (bilateral with stepping, SL with lateral step-down)
 - o Advance core stabilization exercises- SL, Bosu[®], Airex[®], external perturbation, dynamic/rhythmic stabilization exercises, combination of UE and LE with unstable surface and external perturbations, etc.
 - o Isokinetic exercises at (90° - 40°) ($120^{\circ}/s$ - $240^{\circ}/s$), if available
 - o Walking program- 10 minutes minimum, increasing by 10 minutes per week

Advanced Activity Phase:

Criteria for entering this Phase:

- 1- Full knee AROM with no pain or effusion
- 2- Quadriceps Strength & Girth $>80\%$ of the non-injured leg, knee flexor-extensor ratio of 70% to 75%, Hamstring Strength $>80\%$ of the non-injured leg
- 3- No laxity/instability on clinical exam, or KT-2000 test of 2mm or less compared to the non-injured side
- 4- IKDC (use for concomitant injuries)- score of 80% or higher, or the KOOS (use with ACL alone) score of ≥ 80
- 5- Score of "Good " on the LESS to be able to begin Plyometric Training Protocol

Goals for this phase:

- Normalize lower extremity strength, and increase muscle power and endurance
- Maintain/Gain Hamstring & Quadriceps Strength and Girth of 80% or greater
- Continue to improve neuromuscular control
- KOS-sports score of $> 70\%$
- Patient is tested with the Landing Error Scoring System (LESS): **See Attached Sheets: - Test at Patient's individual max vertical jump height, not standard measure on the LESS- may be too high or too low for accurate individual jump assessment score, no sooner than 10 weeks post-op**
 - o Excellent Score is ≤ 4
 - o Good Score is >4 and ≤ 5

- Moderate Score is >5 and ≤ 6
- Poor Score is >6
- Get a baseline FMS® and Y-Balance Assessment™ LE score no sooner than 10-12 weeks Post-op (UE can be testing earlier)
- FMS® (goal of $\geq 14/21$ points with no 0/3, 1/3, or asymmetries) and Y-Balance™ score (goal statistically symmetrical to non-injured leg)
- Perform hop testing $\geq 16-18$ weeks post-op: **See Attached Sheets**
 - 2 practice trials, 2 timed/measured trials; average injured to non-injured
 - 1- Single-leg hop for distance
 - 2- Triple hop for distance
 - 3- Single-Leg Crossover triple hop
 - 4- 6-Metered timed hop

Treatments:

- Exercises: (**Continue/progress all exercises from previous phases as tol./need.**)
 - Advanced core & hip stabilization exercises
 - Begin Plyometric Training Protocol at $\geq 16-18$ weeks * **If LSI $\geq 90\%$ of uninjured leg, with hop tests-** May be delayed longer if allograft reconstruction- patient healing and graft fixation dependant
 - Isokinetic exercises ($180^\circ/s$), if available
 - CKC exercises may be progressed to $75^\circ-90^\circ$ of flexion
 - Eccentric exercises for all lower extremity muscles

Functional Assessments:

- FMS® and Y-Balance Assessment™ **at beginning and end of this phase**
- Repeat the 4 Hop Tests from the previous Phase **at beginning and end of phase**

Core Testing: - **At the beginning of this phase, re-assess any deficits at the end of the phase**

- Segmental Multifidus Test
- Trunk Curl Up Test
- Double-Leg Lowering Test
- Side Bridge Test
- Prone Bridge Test
- Supine Single-Leg Bridge Test
- Extensor Endurance Test

Return to Activity Phase:

Criteria for entering this Phase:

- 1- Full ROM
- 2- A score of $\geq 14/21$ on the FMS® Assessment Screen, with **No** 0/3= pain on any of the 7 fundamental movement patterns
- 3- No statistical asymmetries and on the Y-Balance Assessment™
- 4- Hop Tests (90% or higher compared to non-injured leg)
- 5- Limb Symmetry Index (LSI) of 90% or greater on hop tests
- 6- IKDC Score $\geq 85\%$, or the KOOS score of ≥ 85
- 7- KOS-Sports Score 90% or greater
- 8- Cincinnati Knee Rating System score of 290 points or higher
- 9- No change in knee laxity (clinical exam or ≤ 2 mm on KT 2000 test)
- 10- Isokinetic testing: (if available)
 - i. Quadriceps (80% or greater) compared to non-injured leg
 - ii. Hamstring (85%-100%) compared to non-injured leg
 - iii. Hamstring-Quadriceps Ratio (70% or greater)

Goals or this Phase:

- Achieve maximal strength and endurance
- Normalize neuromuscular control
- Progress to skill training
- Gradually return to sport specific training

Treatments:

Exercises:

- Continue strengthening exercises
- Continue/Advance core training exercises
- Continue Neuromuscular control exercises
- Continue plyometric exercises
 - o Begin walk→run protocol, toward the end of phase, when patient is able to perform a controlled single leg squat (injured leg) to 60° of knee flexion- **see attached sheets**
 - Begin agility and skill training exercises:
 - o Agility exercises- side shuffling, cariocas, zigzags- *** If cleared by MD**
 - o Begin sudden start and stops, figure-8's, 45° and 90° cutting drills, box jumps (progressing & varying heights up to 20cm)- *** If cleared by MD**

Functional Assessment:

- FMS® and Y-Balance Assessment™ **at the end of this phase**
- Repeat the 4 Hop Tests from the previous Phase, Plus Hop-to-Stop Test **at the end of this phase**

Functional Testing: for progression to sport-specific training- (no sooner than 28-45 Weeks) -*Can be perform over multiple days- See Attached Sheets

Strength and Power Testing:

- Single-Leg Squat Test/ Single-Leg Squat Test
- Vertical Jump Test
- Figure-8 Hop Test
- Up-Down Test
- Hexagon Test (DL), Modified Hexagon Hop Test (SL)

Speed, Agility, and Quickness Testing:

- T-Test, or Modified Agility T-Test (MAT)
- Three-Cone Drill Test
- Slalom Test
- Backward Movement Agility Test
- Zigzag Run Test
- Lower Extremity Functional Test (LEFT)

Core Testing:

- Segmental Multifidus Test
- Trunk Curl Up Test
- Double-Leg Lowering Test
- Side Bridge Test
- Prone Bridge Test
- Supine Single-Leg Bridge Test
- Extensor Endurance Test

Function and Balance Testing:

- FMS® Assessment Screen
- Y-Balance Assessment™
- Vail Sports Test™

***Return-to-Sport Protocol- See Specific Sport Return-to-Sport Protocols**

Criteria for Return-to-Sport Specific Protocols:

- A score of $\geq 16/21$ on the FMS® Assessment Screen, with no pain or asymmetries, especially with Inline-lunge, and 2 or greater on all 7 fundamental movement patterns
- Y-Balance Test™- No asymmetries and passing score when compared to peers with data analysis through the Move2Perform software= a composite score as close to 100 as possible
- Limb Symmetry Index (LSI) of $\geq 95\%$ on hop tests
- LEFT Score: ≤ 117 seconds (females), ≥ 105 seconds (males)
- Vail Sport Test™: $\geq 46/54$ points
- Isokinetic testing: (if available)
 - o Quadriceps (90% or greater) compared to non-injured leg
 - o Hamstring (100%-110%) compared to non-injured leg
 - o Hamstring-Quadriceps Ratio (80% or greater)
- IKDC score of $\geq 85\%$, or KOOS score of ≥ 90
- Knee Outcome Survey-Sports Activities Scale (KOS-SAS): $\geq 95\%$
- Cincinnati Knee Rating System: Score ≥ 350
- SL Hop tests $\geq 100\%$ compared to non-injured leg
- No discomfort or swelling, and passing/statistically equal to normative values (if available) with above Functional Tests

References

- Adams D, Logerstedt D, Hunter-Giordano A, Axe MJ, Snyder-Mackler L. Current Concepts for Anterior Cruciate Ligament Reconstruction: A Criterion-Based Rehabilitation Progression, *J Orthop Sports Phys Ther.* 2012; 7:601-614.
- Angelozzi et al. Rate of Force Development as an Adjunctive Measurement for Return-to-Sport Decisions After Anterior Cruciate Ligament Reconstruction. *J Orthop Sports Phys Ther.* 2012; 9:772-780.
- Beynon et al. Accelerated Versus Nonaccelerated Rehabilitation After Anterior Cruciate Ligament Reconstruction: A prospective, Randomized, Double-Blind Investigation Evaluating Knee Joint Laxity Using Roentgen Stereophotogrammetric Analysis. *Am J Sports Med.* 2011; 12:2536-2548.
- Brumitt et al. Lower Extremity Functional Tests and Risk of Injury in Division II Collegiate Athletes. *Inter J Sports Phys Ther.* 2013; 8(3):216-227.
- Christensen, JC, Goldfine, LR, West, HS. The Effects of Early Aggressive Rehabilitation on Outcomes After Anterior Cruciate Ligament Reconstruction Using Autologous Hamstring Tendon: A Randomized Clinical Trial. *Journal of Sports Rehabilitation.* 2013; 22:191-201.
- Cook, Gray. *Movement: Functional Movement Systems* (book). 2010; ON Target Publications.
- Davies GJ, Zillmer DA. Functional Progression of a Patient Through a Rehabilitation Program. *Ortho Phys Ther Clin N Am.* 2000; 9(2):103-117.
- Dunn et al. Which Preoperative Factors, Including Bone Bruise, are Associated with Knee Pain/Symptoms at Index Anterior Cruciate Ligament Reconstruction (ACLR)? A Multicenter Orthopaedic Outcomes Network (MOON) ACLR Cohort Study. *Am J Sports Med.* 2010; 9:1778-1787.
- Eitzen I, Mosksnes H, Snyder-Mackler L, Risberg MA. A Progressive 5-Week Exercise Therapy Program Leads to Significant Improvement in Knee Function Early After Anterior Cruciate Ligament Injury, *J Ortho Sports Phys Ther.* 2010; 11:705-721.
- Garrison et al. The Reliability of the Vail Sport Test™ as a Measure of Physical Performance Following Anterior Cruciate Ligament Reconstruction. *Inter J Sports Phys Ther.* 2012; 7(1):20-30.
- Goulbourne, Shaun. *Sports-Related Injuries in the Young Athlete* (book from training course). 2011; Cross Country Education, LLC.
- Hewett TE, Stroupe AL, Noyes FR. Plyometric Training in Female Athletes. Decreased Impact forces and Increased Hamstring Torques. *Am J Sports Med.* 1996; 2:765-773.
- Lentz et al. Return to Preinjury Sports Participation Following Anterior Cruciate Ligament Reconstruction: Contributions of Demographic, Knee Impairments, and Self-Report Measures. *J Orthop Sports Phys Ther.* 2012; 11:893-901.
- Myer GD, Paterno MV, Ford KR, Quatman CE, Hewett TE. Rehabilitation After Anterior Cruciate Ligament Reconstruction: Criteria-Based Progression through the Return-to-Sport Phase. *J Orthop Sports Phys Ther.* 2006; 6:385-402.

Narducci E, Waltz A, Gorski K, Leppla L, Donaldson M. The Clinical Utility of Functional Performance Tests Within One-Year Post-ACL Reconstruction: A Systematic Review. *Inter J Sports Phys Ther.* 2011; 4:333-342.

Padua et al. The Landing Error Scoring System (LESS) is a Valid and Reliable Clinical Assessment Tool of Jump-Landing Biomechanics: The Jump-ACL Study, *Am J Sports Med.* 2009; 10:1996-1002.

Paterno et al. Biomechanical Measures During Landing and Postural Stability Predict Second Anterior Cruciate Ligament Injury After Anterior Cruciate Ligament Reconstruction and Return to Sport, *Am J Sports Med.* 2010; 10:1968-1978.

Paterno MV, Rauh MJ, Schmitt LC, Ford KR, Hewett TE. Incidence of Contralateral and Ipsilateral Anterior Cruciate Ligament (ACL) injury After Primary ACL Reconstruction and Return to Sport. *Clin J Sport Med.* 2012; 2: 116-121.

Reiman, Michael P., Manske, Robert C. *Functional Testing in Human Performance: 139 tests for sport, fitness, and Occupational Settings* (book). 2009. Human Kinetics.

Ross MD, Langford B, Whelan PJ. Test-retest Reliability of 4 Single-Leg Horizontal Hop Tests. *J Strength Cond Res.* 2002; 4:617-622.

Schmitt LC, Paterno MV, Hewett TE. The Impact of Quadriceps Femoris Strength Asymmetry on Functional Performance at Return to Sport Following Anterior Cruciate Ligament Reconstruction. *J Orthop Sports Phys Ther.* 2012; 9:750-759.

Spindler et al. Prognosis and Predictors of ACL Reconstructions using the MOON Cohort: A Model for Comparative Effectiveness Studies. *J Ortho Res.* 2013; 1:2-9.

Stearns KM, Pollard CD. Abnormal Frontal Plane Knee Mechanics During Sidestep Cutting in Female Soccer Athletes After Anterior Cruciate Ligament Reconstruction and Return to Sport. *Am J Sports Med.* 2013; 4:918-923.

Van Ginckel A, Vedonk P, Victor J, Witvrouw E. Cartilage Status in Relation to Return to Sports After Anterior Cruciate Ligament Reconstruction. *Am J Sports Med.* 2013; 3:550-559.

Wilk KE, Macrina LC, Cain EL, Dugas JR, Andrews JR. Recent Advances in the Rehabilitation of Anterior Cruciate Ligament Injuries. *J Orthop Sports Phys Ther.* 2012; 3: 153-171.

Yabroudi MA, Irrgang JJ. Rehabilitation and Return to Play After Anatomic Anterior Cruciate Ligament Reconstruction. *Clin Sports Med.* 2013 165-175.