



ORTHOPEDIC ASSOCIATES of HARTFORD

Chris Lena MD, James Alvarez PAC

Arthroscopic and Reconstructive Surgery of the Shoulder and Knee Sports Medicine

MEA's Karen Smith, Jackie Zuidema, Annmarie Fiore

Tel: (860) 549-8249 - FAX: (860) 244-8813

www.oahct.com

Arthroscopic Chondroplasty/Subchondroplasty

- 1. ACTIVITIES:** For 1-2 weeks after surgery, you will be asked to maintain 50% weight-bearing on the operative leg using crutches. The use of crutches can typically be reduced as tolerated. You are encouraged to try to walk as smoothly as possible. Do not do any strenuous activity such as running and jumping until I clear you for this. You may start bending your knee as tolerated, the sooner the better.
- 2. BANDAGES:** Your bandage may be removed 2 days following surgery. The knee should then be re-wrapped with only the elastic bandage for about 3-4 days or until swelling is gone. **DO NOT** wrap the elastic bandage too tightly, or it will act like a tourniquet and cause ankle swelling.
- 3. MEDICATIONS:** A prescription will be provided to help relieve pain. Please use this medication as directed. This medication is strong, and should not be taken with alcohol or other pain medications, and may cause drowsiness. Exercise good judgment in its use. You may also try over the counter pain medications such as Aleve (Naprosyn) or Advil (Ibuprofen). Take as directed unless there are contraindications. **Take one aspirin (325 mg) per day.**
- 4. SHOWER:** You may shower after 48 hours. Do not take a bath or submerge the knee under water for 7 days. If any drainage is present, do not get the knee wet and please call the office.
- 5. STITCHES:** There are stitches in the skin. The small skin incisions are held together with sutures, and if any problem is noted with the incision, please call the office.
- 6. EXERCISES:** The thigh muscles will shrink in size and strength quite rapidly unless they are exercised. Simple exercises should be started as soon as possible. The best exercises are as follows:
 - Straighten your knee as much as possible and clench the thigh muscles tightly. Hold the muscles clenched tight for 5 seconds, and then relax. Repeat this exercise 10-20 times every 30-60 minutes. You should try to do at least 100 tightenings per day to keep the tone and strength in the muscle.
 - Start bending your knee the day after surgery and increase the bending until full motion has returned.
 - A stationary bicycle may be used once the knee bends to 90 degrees.
 - The need for physical therapy will be discussed at your follow-up visit
- 7. WORK ACTIVITY:** Most people are able to return to either sedentary or restricted work activity within 1-3 days of their operative procedure. Generally, full normal stressful activities and sports are resumed after 4-6 weeks. During this time frame, more stressful activities may be engaged as tolerated.
- 8. COMFORT:** Pain and swelling after arthroscopy is to be expected. This should subside after 2-4 days. If the following develop and persist after 24 hours, please call the office:
 - Fever over 101 degrees Fahrenheit that is not relieved with the use of antipyretics (such as Tylenol, Aspirin or Ibuprofen) should be called to your doctor's attention.
 - Swelling below the knee, in the calf, ankle or foot which does not respond to loosening of the ace wrap, ice and elevation.
 - Increasing pain in the knee or calf.
 - Discharge or drainage from the knee which continues greater than 24 hours.
- 9. FOLLOW-UP:** Please call the office the day after your operation and make an appointment for 7-10 days after your procedure. You will be checked, your procedure discussed and your rehabilitation will be planned.

KNEE MENISCECTOMY/CHONDROPLASTY/SUBCHONDROPLASTY

POSTOPERATIVE INSTRUCTIONS

You've had arthroscopic surgery today. Your knee joint was evaluated thoroughly.

I found in your knee:

- | | | |
|--|--|--|
| <input type="checkbox"/> Medial meniscal tear | <input type="checkbox"/> Arthritis | <input type="checkbox"/> Ligament Damage |
| <input type="checkbox"/> Lateral meniscal tear
trochlea | <input type="checkbox"/> Cartilage Damage Patella – medial femoral – lateral femoral – | |
| <input type="checkbox"/> Loose bodies | <input type="checkbox"/> Osteophytes | <input type="checkbox"/> Plica |
-

I was able to:

- | | | | |
|---|---|--|---|
| <input type="checkbox"/> Debride meniscus | <input type="checkbox"/> Clean Arthritis | <input type="checkbox"/> Remove loose bodies | <input type="checkbox"/> Remove plica |
| <input type="checkbox"/> Repair meniscus | <input type="checkbox"/> Shave cartilage damage | <input type="checkbox"/> Remove osteophytes | <input type="checkbox"/> Subchondroplasty |
-

Until I see you, you should:

- Use crutches Try to walk normally Straighten and bend knee
- Start physical therapy (a referral has been given)

You should change your dressing in 2 days. Cover the portals with band aids. We will remove all stitches at your visit. You may shower at 48 hours but do not soak the incisions.

Pain is expected for several weeks. Use your medication for severe pain. Take Aleve or Advil as directed as needed (if not allergic, or no ulcer history). Call if fever greater than 101 °F and not relieved with Tylenol, or if infection is noted. Call 549-8249 for office visit **7-10** days after surgery. If you want to return to work before your next visit, call the office.

